

Triplication of $SU(5)$ monopoles

Levon Pogosian

*Theoretical Physics, The Blackett Laboratory, Imperial College,
Prince Consort Road, London SW7 2BZ, United Kingdom.*

Danièle A. Steer

Laboratoire de Physique Théorique, Bât. 210, Université Paris XI, 91405 Orsay Cedex, France.

Tanmay Vachaspati

*Department of Physics, Case Western Reserve University,
10900 Euclid Avenue, Cleveland, OH 44106-7079, USA.*

We investigate all spherically symmetric fundamental monopole solutions with fixed topological charge in the $\tilde{S}U(5) \rightarrow [\tilde{S}U(3) \times \tilde{S}U(2) \times \tilde{U}(1)]/\tilde{Z}_3 \times \tilde{Z}_2$ symmetry breaking. We find that there is three-fold replication of the monopoles. The three copies correspond to bound states with “monopole clouds” arising from the non-Abelian nature of the $\tilde{S}U(3)$ factor. The triplication of monopoles could help us understand the observed family structure of standard model particles.

PACS numbers: 98.80.Cq

Magnetic monopoles are of diverse and fundamental interest since they are predicted by all grand unified theories and embody a rich mathematical structure. They are also central to discussions of non-perturbative field theory and may provide a new perspective on particle physics phenomenology.

The topological properties of any field theory leads to a classification of the topological defect solutions it contains. However, there are interesting and important properties of field theories that are missed by the topological classification. For example, a more detailed study of domain walls in an $SU(5) \times Z_2$ model reveals three different domain wall solutions, all having the same topological charge [1, 2]. Guided by the domain wall example, we turn our attention to magnetic monopole solutions in

$$\tilde{S}U(5) \rightarrow [\tilde{S}U(3) \times \tilde{S}U(2) \times \tilde{U}(1)]/\tilde{Z}_3 \times \tilde{Z}_2 \quad (1)$$

(The tildes indicate that the groups should not be confused with the symmetry groups of the standard model, which will be written without tildes.) This symmetry breaking is of interest because it is the basis of current grand unification models. More importantly for us is the fact that the topological charges of (non-BPS) magnetic monopole solutions in this model are in one to one correspondence with the electric charges of the standard model leptons and quarks [3]. If duality holds, then any structure of the $\tilde{S}U(5)$ monopole solutions should be reflected in the standard model particles and vice versa. In particular we ask: can the family structure of standard model particles be present in the monopole solutions? (A different approach to the family structure was taken in [4].)

Our approach to solving the problem is to construct all spherically symmetric monopole solutions with unit winding and fixed charge orientation in the model. Here

we are not interested in continuous degeneracies of the solutions that are given by the zero modes. Instead we are looking for features in the space of solutions, particularly for non-BPS monopoles, that might point to a replication in the quantum theory of monopoles. In our effort to determine all such solutions, we rely on the construction of Wilkinson and Goldhaber (WG) [5], though with a novel twist that we detail below. We are led to three possible solutions that are then found numerically. Two of these have the strange feature that they are not localized and hence need careful interpretation. Such a situation was analyzed in detail by Weinberg and collaborators [6, 7] in the BPS case and, following their discussion, we interpret each of our two non-localized solutions as a “monopole plus cloud”. A further symmetry breaking of the $\tilde{S}U(3)$ group in eq. (1) to $\tilde{U}(1)_1 \times \tilde{U}(1)_2$ would condense the clouds into localized objects. Now the interpretation of the three monopoles as distinct objects is very clear since they all have different masses.

We briefly discuss the WG [5] recipe for constructing spherically symmetric magnetic monopole solutions. (This scheme has earlier been applied to the $\tilde{S}U(5)$ case in [8, 9] but in a different context.) We start by fixing the vacuum expectation value (VEV) of the $\tilde{S}U(5)$ adjoint scalar field, Φ , along the z -axis:

$$\Phi(z \rightarrow \infty, \hat{\mathbf{z}}) \equiv \Phi_0 = \frac{\eta}{2\sqrt{15}} \text{diag}(2, 2, 2, -3, -3) \quad (2)$$

where η is the magnitude of the VEV. We also fix the magnetic charge, Q , of the magnetic monopole solution:

$$Q = -\text{diag}(0, 0, \sigma_3, 0)/2. \quad (3)$$

(σ_a , $a = 1, 2, 3$ denote the Pauli spin matrices.) Then the asymptotic magnetic field along the z -axis is:

$$e\mathbf{B} = \frac{Q}{r^2} \hat{\mathbf{r}}. \quad (4)$$

where e is the gauge coupling. This monopole corresponds to the left-handed quark doublet in the dual standard model [3].

We would like to find all spherically symmetric solutions corresponding to monopoles with charge Q given in eq. (3). The key idea of the WG recipe is to start by working in the singular Abelian gauge - the gauge in which monopoles are point objects. In this gauge Φ is fixed, $\Phi = \Phi_0$, and the gauge fields are those of a Dirac monopole. It is relatively straightforward to write down all possible point-monopole solutions given the symmetries of the system. WG then prescribe how to construct a gauge transformation that would transform each of these point-monopoles into spherically symmetric finite-energy monopole solutions.

As described by WG [5], though changing the sequence of their reasoning, we must find all $SU(2)$ generators I_a such that

$$[I_a, Q] = 0, \quad [I_a, \Phi_0] = 0, \quad a = 1, 2, 3. \quad (5)$$

The choice of I_a fixes the third generator, T_3 , of the $SU(2)$ that defines the spherical symmetry of the solution:

$$T_3 = I_3 - Q. \quad (6)$$

Note that $[Q, \Phi_0] = 0$ and therefore $[T_3, \Phi_0] = 0$. Once T_3 is known, we can find the ladder matrices $T_{\pm}$ satisfying:

$$[T_3, T_{\pm}] = \pm T_{\pm}, \quad (7)$$

and hence the T_1 and T_2 generators via $T_1 = (T_+ + T_-)/2$ and $T_2 = (T_+ - T_-)/2i$. Note that if T_3 has degenerate eigenvalues, there can be several choices for the $T_{\pm}$. The asymptotic scalar fields are written as:

$$\Phi(\hat{\mathbf{r}}) = \Omega(\hat{\mathbf{r}})\Phi_0\Omega^{-1}(\hat{\mathbf{r}}) \quad (8)$$

where $\hat{\mathbf{r}} = (\sin\theta\cos\phi, \sin\theta\sin\phi, \cos\theta)$, and $\Omega(\hat{\mathbf{r}}) = e^{-i\phi T_3}e^{-i\theta T_2}e^{i\phi T_3}$. On the other hand, the asymptotic gauge fields are given by

$$e\mathbf{A} = [\mathbf{I}(\hat{\mathbf{r}}) - \mathbf{T}] \times \frac{\hat{\mathbf{r}}}{r} \quad (9)$$

where $\mathbf{I}(\hat{\mathbf{r}}) = \Omega(\hat{\mathbf{r}})\omega^{-1}(\hat{\mathbf{r}})\mathbf{I}\omega(\hat{\mathbf{r}})\Omega^{-1}(\hat{\mathbf{r}})$ with $\omega(\hat{\mathbf{r}}) = e^{-i\phi I_3}e^{-i\theta I_2}e^{i\phi I_3}$. Note that where I_a is written without any argument, it refers to $I_a(\hat{\mathbf{z}})$.

Finally, we need to introduce profile functions to get the radial dependence of the fields [5]. This leads to the following ansatz for the scalar field along the z -axis:

$$\begin{aligned} r\Phi(r) &= r \text{diag}(\Phi_{11}, \Phi_{22}, \Phi_{33}, \Phi_{44}, \Phi_{55}) \\ &= h_1(r)\lambda_3 + h_2(r)\lambda_8 + h_3(r)\tau_3 + h_4(r)Y \end{aligned} \quad (10)$$

where

$$\begin{aligned} \lambda_3 &= \frac{1}{2}\text{diag}(\sigma_3, 0, 0, 0), \quad \lambda_8 = \frac{1}{2\sqrt{3}}\text{diag}(1, 1, -2, 0, 0), \\ \tau_3 &= \frac{1}{2}\text{diag}(0, 0, 0, \sigma_3), \quad Y = \frac{1}{2\sqrt{15}}\text{diag}(2, 2, 2, -3, -3). \end{aligned}$$

The magnetic field is given in terms of complex functions, $v_{ij}(r)$, $i, j = 1, \dots, 5$ but only four of them will be non-zero: the function v_{lk} is non-trivial only if there is a possible choice of T_+ such that $[T_+]_{lk}$ is non-zero [5]. Next we define

$$[M_1(r)]_{ij} = \frac{v_{ij} + v_{ji}^*}{2}, \quad [M_2(r)]_{ij} = \frac{v_{ij} - v_{ji}^*}{2i} \quad (11)$$

from which the gauge field is:

$$e\mathbf{A}(r) = [\mathbf{M}(r) - \mathbf{T}] \times \frac{\hat{\mathbf{r}}}{r}. \quad (12)$$

The energy functional is

$$E = \int d^3x [\text{Tr}\mathbf{B}^2 + \text{Tr}(\mathbf{D}\Phi)^2 + V(\Phi)] \quad (13)$$

with

$$V(\Phi) = -m^2\text{Tr}(\Phi^2) + h(\text{Tr}(\Phi^2))^2 + \lambda\text{Tr}(\Phi^4) - \frac{m^4}{4\lambda'}, \quad (14)$$

where $\lambda' = h + 7\lambda/30$ and $\eta = m/\sqrt{\lambda'}$ (see Eq. (2)).

Different functions v_{ij} will be non-trivial, depending on the choice of $\mathbf{I}$ satisfying eq. (5). The only possibilities for I_3 are: $I_3 = 0$ or $I_3 = \text{diag}(\pm\sigma_3, 0, 0, 0)/2$. The former choice gives the conventional monopole [8, 9], while it turns out that the latter can lead to new monopoles solutions in addition to the conventional one. With $\mathbf{I} = \text{diag}(+\sigma, 0, 0, 0)/2$, the four non-trivial gauge profile functions are: v_{12} , v_{14} , v_{32} and v_{34} corresponding to the 4 different non-zero entries in the two possible constructions of T_+ . We will write the equations of motion only for this choice of $\mathbf{I}$.

Inserting the ansatz for the fields into the energy functional and then extremizing, leads to the equations of motion for all spherically symmetric monopole solutions:

$$\begin{aligned} r^2 v_{12}'' &= v_{12}(v_{12}^2 + v_{14}^2 + v_{32}^2 + e^2 H_{12}^2 - 1) - v_{14}v_{32}v_{34}, \\ r^2 v_{14}'' &= v_{14}(v_{14}^2 + v_{12}^2 + v_{34}^2 + e^2 H_{14}^2 - 1) - v_{12}v_{32}v_{34}, \\ r^2 v_{32}'' &= v_{32}(v_{32}^2 + v_{12}^2 + v_{34}^2 + e^2 H_{23}^2 - 1) - v_{12}v_{14}v_{34}, \\ r^2 v_{34}'' &= v_{34}(v_{34}^2 + v_{14}^2 + v_{32}^2 + e^2 H_{34}^2 - 1) - v_{12}v_{14}v_{32}, \end{aligned}$$

$$\begin{aligned} r^2 h_1'' &= 2H_{12}v_{12}^2 + H_{14}v_{14}^2 - H_{23}v_{32}^2 + U_1, \\ r^2 h_2'' &= \frac{1}{\sqrt{3}}H_{14}v_{14}^2 + \sqrt{3}H_{23}v_{32}^2 - \frac{2}{\sqrt{3}}H_{34}v_{34}^2 + U_2, \\ r^2 h_3'' &= -H_{14}v_{14}^2 - H_{34}v_{34}^2 + U_3, \\ r^2 h_4'' &= \frac{\sqrt{15}}{3}H_{14}v_{14}^2 + \frac{\sqrt{15}}{3}H_{34}v_{34}^2 + U_4, \end{aligned}$$

where primes denote derivatives with respect to r , U_i are derivatives of the potential that we will not explicitly write out, and $H_{ij} \equiv r(\Phi_{ii} - \Phi_{jj})$. In the above equations we have already chosen the phases, θ_{ij} , of the complex functions v_{ij} such that the energy is minimized. These phases are constant in space and satisfy $\theta_{12} + \theta_{34} - \theta_{14} -$

$\theta_{32} = \pi$, resulting in the minus sign in front of the last term in the equations for the v_{ij} 's.

The boundary conditions on the Higgs field at $r = \infty$ are determined by requiring that $\Phi = \Phi_0$ asymptotically and those on the gauge field by requiring that eq. (12) goes over into eq. (9) at infinity. Then,

$$\begin{aligned} v_{12}(\infty) &= 1, & v_{14}(\infty) &= v_{32}(\infty) = v_{34}(\infty) = 0 \\ h_1(\infty) &= 0, & h'_2(\infty) &= h'_3(\infty) = 0, & h'_4(\infty) &= \eta. \end{aligned}$$

The boundary conditions on the Higgs field at the origin are determined by requiring non-singular behavior:

$$h_i(0) = 0, \quad i = 1, 2, 3, 4, \quad (15)$$

and the functions v_{ij} at the origin are as yet unspecified. This is where we encounter the freedom that ultimately leads to three different monopole solutions. As this is the crucial new twist in the recipe, we describe it in some detail.

Going back to the WG recipe, notice that if I_3 is $\text{diag}(\sigma_3, 0, 0, 0)/2$ then $T_3 = \text{diag}(1, -1, 1, -1, 0)/2$. This leads to two possible choices for $T_{\pm}$. The first is when T_+ is non-trivial in the 1-2 and 3-4 blocks. The second choice is when T_+ is non-trivial in the 1-4 and 3-2 blocks. Requiring that the gauge field in eq. (12) be non-singular at the origin fixes the values of functions v_{ij} at $r = 0$. Then, for the case where T_+ is in the 1-2 and 3-4 blocks we have the boundary conditions:

$$v_{12}(0) = 1 = v_{34}(0); \quad v_{14}(0) = 0 = v_{32}(0). \quad (16)$$

When T_+ is chosen in the 1-4 and 3-2 blocks:

$$v_{14}(0) = 1 = v_{32}(0); \quad v_{12}(0) = 0 = v_{34}(0). \quad (17)$$

Similarly, we can also choose $I_3 = \text{diag}(-\sigma_3, 0, 0, 0)/2$ and then $T_3 = \text{diag}(-1, 1, 1, -1, 0)/2$. This leads to T_+ lying either in the 3-1 and 2-4 blocks, or in the 2-1 and 3-4 blocks. The non-trivial gauge profile functions are now: v_{21} , v_{24} , v_{31} and v_{34} [5]. The second choice for T_+ can be shown to lead to a solution that has scalar and magnetic fields that are identical to those in the first case (eq. (16)). The boundary conditions on the profile functions for the first choice are determined by requiring that the gauge field remain non-singular at the origin:

$$v_{31}(0) = 1 = v_{24}(0); \quad v_{21}(0) = 0 = v_{34}(0). \quad (18)$$

In addition, the boundary conditions at infinity are $v_{21}(\infty) = 1$ and $v_{ij}(\infty) = 0$ for all $(i, j) \neq (2, 1)$. This is the third possible monopole solution.

Now our system is complete – we have the differential equations and the boundary conditions. We solve the equations numerically with the various boundary conditions. The first case (eq. (16)) leads to the conventional magnetic monopole which is also discussed in Refs. [8, 9] (The conventional monopole is usually constructed by setting $I_3 = 0$ and repeating the above procedure). The second and third set of boundary conditions give rise to




FIG. 1: Profile functions $v_{ij}(r)$ and $h_i(r)$ for $e = 1$, $\lambda = 0.1$, $h/\lambda = -0.2$ and $\eta = 1$ (see Eq. (14) for definitions). The upper panel shows plots of v_{12} (solid), v_{14} (dot), v_{32} (short dash) and v_{34} (long dash). The function v_{34} is not identically zero but vanishes in the limit of an infinite lattice size. The lower panel shows plots of h_1/r (solid), h_2/r (dot), h_3/r (short dash) and h_4/r (long dash). Functions v_{12} and v_{32} approach their boundary values linearly.

two new solutions. The profile functions for the second case are shown in Fig. 1. The third monopole solution can be obtained from the second by switching the 1 and 2 rows and columns of the fields and simply amounts to re-labeling the profile functions. Note that, while the conventional monopole obtained using the boundary conditions given in eq. (16) is the same as the one obtained by choosing $I_3 = 0$, there will be a difference between the two if one breaks the $\tilde{S}U(3)$ symmetry. This will become relevant later when we consider the case of $\tilde{S}U(3)$ breaking down to $\tilde{U}(1)_1 \times \tilde{U}(1)_2$. With $I_3 = 0$, the gauge profile functions must obey the boundary conditions $v_{ij}(\infty) = 0$, $v_{34}(0) = 1$ with the rest of v_{ij} vanishing at $r = 0$.

The plot in Fig. 1 shows some unusual features. While most of the profile functions show “good” asymptotic behavior, the functions v_{12} and v_{32} approach their boundary values linearly. This behavior can be confirmed by analytically expanding the equations of motion. It is also physically reasonable since the functions H_{12} and H_{23} appearing on the right-hand sides of the equations of motion do not acquire a VEV. Therefore the functions v_{12} and v_{32} correspond to massless gauge degrees of freedom. The energy of the solution approaches the energy of the conventional monopole as the lattice gets bigger.

We now interpret this rather peculiar situation following the discussion of BPS magnetic monopoles in the presence of unbroken non-Abelian symmetries by Wein-

berg and collaborators [6, 7]. The idea is to consider the case of maximal symmetry breaking first and then to decrease the VEVs such that, in the limit of zero VEV, the original non-Abelian symmetry is restored. In our case, we only need to consider the breaking of the $\tilde{SU}(3)$ group further down to $\tilde{U}(1)_1 \times \tilde{U}(1)_2$. In the numerical implementation described above, the symmetry breaking is achieved by giving a VEV to h_1 and setting $v_{ij}(\infty) = 0$. Working, for simplicity, in the BPS limit (*i.e.* $V(\Phi) = 0$), we now let $h_1 = \eta_1 r$ in the asymptotic region. When $\eta_1 \neq 0$, the symmetry breaking is:

$$\tilde{SU}(5) \rightarrow [\tilde{U}(1)_1 \times \tilde{U}(1)_2 \times \tilde{SU}(2) \times \tilde{U}(1)] / \tilde{Z}_3 \times \tilde{Z}_2 \quad (19)$$

As η_1 is tuned to zero, we obtain the symmetry breaking pattern in eq. (1). Now the equations of motion are solved again with the appropriate boundary conditions.

Just as we might expect, with $\eta_1 \neq 0$, all the three monopoles are well-localized and have different masses. In the case of the conventional monopole, obtained using $I_3 = 0$, the $\tilde{U}(1)_1$ magnetic charge of the solution is zero and the topological charge of the monopole is

$$Q = \frac{1}{2} \text{diag}(0, 0, -1, 1, 0) . \quad (20)$$

The topological charge for the monopole in the second case (eq. (17)) after the breaking of the $\tilde{SU}(3)$ symmetry is

$$Q' = Q + \frac{1}{2} \text{diag}(-1, 1, 0, 0, 0) , \quad (21)$$

where the second term is the charge of the $\tilde{U}(1)_1 \times \tilde{U}(1)_2$ monopole which only condenses when $\tilde{SU}(3)$ is broken. The topological charge of the monopole in the third case is

$$Q'' = Q + \frac{1}{2} \text{diag}(1, -1, 0, 0, 0) . \quad (22)$$

As η_1 is decreased, the $\tilde{U}(1)_1 \times \tilde{U}(1)_2$ monopole will spread, ultimately filling all space and becoming a “cloud”. As in the BPS analysis [7], the energy associated with the cloud goes to zero when we extrapolate to an infinite lattice. With $\eta_1 = 0$, we can explicitly check that monopoles 2 and 3 are gauge equivalent. The gauge equivalence of monopoles 1 and 2 (and, 1 and 3)

is more delicate since the cloud, which is the cause of gauge inequivalence on a finite lattice, gets pushed to infinity in the infinite lattice limit. The two solutions are equivalent but only under large gauge transformations. As described below, this delicate behavior is not relevant for us since we will be interested in the case when $\tilde{SU}(3)$ gets broken.

Therefore, we have in hand a total of $3 \times (3 \times 2) = 18$ gauge equivalent fundamental monopole solutions in the symmetry breaking in eq. (1). The 3×2 factor is due to the 3 permutations of the λ_8 and the 2 permutations of τ_3 , and the factor of 3 refers to the three solutions described above. Classically there is no distinction between these solutions. However, in a dual quantum theory the $3 \times 2 = 6$ solutions are believed to correspond [10] to 3 color ($SU(3)_c$) and 2 isospin ($SU(2)_L$) degrees of freedom [3]. The remaining factor of 3 suggests that there is a 3-fold replication of degrees of freedom in the quantum theory even for fixed color, isospin and hypercharge ($U(1)_Y$).

In the context of the dual standard model, the interpretation of the 3 solutions is clear. The reason is that color confinement in the dual standard model is obtained by further breaking $\tilde{SU}(3)$ to $\tilde{Z}_3$. Then the colored monopoles get connected by $\tilde{Z}_3$ strings in color singlet objects. However, the three solutions are manifestly distinct when $\tilde{SU}(3)$ is broken, since then they have different masses. Hence they necessarily represent distinct particles in the dual (electric) theory.

Higher winding monopoles are built by combining fundamental monopoles and so we expect them to be replicated as well. It would be very interesting to see if the entire monopole spectrum is exactly triplicated. If there are indeed three families of monopole solutions, it would complete the tantalizing correspondence between the magnetic charge spectrum of $\tilde{SU}(5)$ monopoles and the electric charge spectrum of observed particles [3].

TV is grateful to Fred Goldhaber and Erick Weinberg for useful discussions. DS is grateful to the Swiss Foundation Ernst et Lucie Schmidheiny for travel support and to Case Western Reserve University for hospitality. This collaboration was in part possible due to support by the ESF COSLAB network. LP was supported by PPARC and TV by DOE.

[1] T. Vachaspati, Phys. Rev. **D63**, 105010 (2001).
[2] L. Pogosian and T. Vachaspati, Phys. Rev. **D64**, 105023 (2001).
[3] T. Vachaspati, Phys. Rev. Lett. **76**, 188 (1996); H. Liu and T. Vachaspati, Phys. Rev. **D56**, 1300 (1997).
[4] H. Liu, G.D. Starkman and T. Vachaspati, Phys. Rev. Lett. **78**, 1223 (1997).
[5] D. Wilkinson and A.S. Goldhaber, Phys. Rev. **D16**, 1221 (1977).

[6] E.J. Weinberg, Nucl. Phys. **B203**, 445 (1982).
[7] K. Lee, E.J. Weinberg and P. Yi, Phys. Rev. **D54**, 6351 (1996).
[8] C.P. Dokos and T.N. Tomaras, Phys. Rev. **D21**, 2940 (1980).
[9] D.M. Scott, Nucl. Phys. **B171**, 95 (1980).
[10] P. Goddard, J. Nuyts and D.I. Olive, Nucl. Phys. **B125**, 1 (1977).